The U.S. Geological Survey (USGS), Esri, the Group on Earth Observations (GEO), and the American Association of Geographers (AAG) are pleased to present A New Map of Global Islands. This publication describes the development of a new global islands geospatial data resource, and presents maps and a web-based explorer application for visualizing the data. The islands were mapped from satellite imagery, and represent a high spatial resolution, globally comprehensive characterization of over 340,000 islands. The islands range in size from very large continental mainlands to islands as small as half of a soccer field.

The islands and associated shoreline data were developed as a first step towards the mapping and characterization of global ecological coastal units (ECUs). The work to delineate the ECUs follows previous efforts to map global ecological land units (ELUs) and ecological marine units (EMUs). These global ecosystem mapping resources have been commissioned by GEO, a consortium of over 200 nations seeking to advance the use of satellite and in-situ observations of the planet for societal benefit. With this Special Publication, AAG recognizes the work to better characterize the world's islands as a fundamental exploration in physical and ecological geography.



A Special Publication of the American Association of Geographers



-







Major contributors to this publication include:

of GEOGRAPHERS







The American Association of Geographers is a nonprofit scientific and educational society with a membership of almost 12,000 individuals from more than 60 countries. AAG members are geographers and related professionals who work in the public, private, and academic sectors to advance the theory, methods, and practice of geography.

The U.S. Geological Survey (USGS) was created in 1879 as a science agency charged with providing information and understanding to help resolve complex natural resource problems across the nation and around the world. The mission of the USGS is to provide relevant, impartial scientific information to 1) describe and understand the Earth, 2) minimize loss of life and property from natural disasters, 3) manage water, biological, energy, and mineral resources, and 4) enhance and protect our quality of life.

Esri is an international supplier and leader in geographic information system (GIS) software, research and development. Its flagship product, ArcGIS, is the world's most powerful mapping and spatial analytics software, applying The Science of Where to connect everyone, everywhere through a common visual language. This combination of mapping and analytics reveals deeper insight into the world's data, and is enabling over 350,000 organizations worldwide in creating responsible and sustainable solutions to problems, from local to global scales.

The Group on Earth Observations (GEO) is a voluntary, international partnership of governments and scientific and technical organizations collaborating to develop a Global Earth Observation System of Systems (GEOSS). GEO's vision is to realize a future wherein decisions and actions for the benefit of humankind are informed by coordinated, comprehensive and sustained Earth observations and information. GEO BON is GEO's Biodiversity Observation Network, and GEO ECO is an initiative to map and monitor global ecosystems.

Roger Sayre, U.S. Geological Survey, Reston, Virginia Jack Dangermond, Esri, Redlands, California Dawn Wright, Esri, Redlands, California Sean Breyer, Esri, Redlands, California Kevin Butler, Esri, Redlands, California Keith Van Graafeiland, Esri, Redlands, California Charles Frye, Esri, Redlands, California Deniz Karagulle, Esri, Redlands, California Steve Kopp, Esri, Redlands, California Suzanne Noble, U.S. Geological Survey, Denver, Colorado Jill Cress, U.S. Geological Survey, Denver, Colorado Devon Burton, U.S. Geological Survey, Reston, Virginia Madeline Martin, U.S. Geological Survey, Reston, Virginia Jac Steiner, University of Colorado Denver, Denver, Colorado



© American Association of Geographers, 2019 1710 16th Street NW, Washington, DC 20009-3198 • www.aag.org All rights reserved.

Published by the Association of American Geographers in collaboration with the U.S. Geological Survey (USGS), Esri, and the Group on Earth Observations (GEO). The first author is responsible for the choice and the presentation of material contained in this publication, and any opinions expressed do not necessarily reflect the views or policies of the publisher, AAG, or the government of United States. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

ISBN 978-0-89291-284-1

Cover design by Roger Sayre (USGS) and Becky Pendergast (AAG).

Citation: Sayre, R., J. Dangermond, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, C. Frye, D. Karagulle, S. Kopp, S. Noble, J. Cress, D. Burton, M. Martin, and J. Steiner. 2019. A New Map of Global Islands. Washington, DC: American Association of Geographers. 24 pages.

A New Map of Global Islands

Earth, sea and sky — A view of a small island off the coast of Eriskay in the Outer Hebrides, Scotland (Photo by Becky Pendergast)

Foreword

A New Map of Global Islands

Following on the publication of Global Ecological Land Units (2014) and Global Ecological Marine Units (2017), AAG is pleased to present a third booklet in its Special Publication Series on Global Ecosystem Mapping. AAG is collaborating with the U.S. Geological Survey, Esri, and the Group on Earth Observations (GEO) to publish standardized, robust and practical maps of global ecosystems for a variety of research and management applications. The global ecosystem mapping work is commissioned by GEO, a consortium of nations dedicated to advancing the use of earth observations for societal benefit.

This publication maps hundreds of thousands of islands in greater detail than previous efforts. The islands range in size from continental mainlands to tiny islets smaller than a fraction of a square kilometer. The work was done at a very fine spatial resolution, and the shorelines and islands were mapped from hundreds of 2014 Landsat images.

The publication also describes an elegantly simple tool, the Global Island Explorer, which was designed as a window for users of the data. The Global Island Explorer is an online visualization and query tool, which allows anyone with an internet connection to explore any of the world's islands in an easy to use app.

Like the global ecological land and marine units before them, the production of the new Global Islands resource was strongly supported by Esri. Jack Dangermond is to be commended for his appreciation of the importance of mapping and understanding global ecosystems, and for engaging in the work and hosting the content. The USGS is also to be commended for their leadership in the global ecosystems mapping initiative, and Roger Sayre's unwavering dedication to advancing the work. The AAG is pleased to contribute to this ongoing global ecosystem mapping work. We recognize and celebrate this important exploration in applied physical geography.

Douglas Richardson Executive Director American Association of Geographers

Abstract	
Introduction	
Geographic Data on Global Islands	6
Islands and Biodiversity	
The GEO (Group on Earth Observations) Global Ecosystem Mapping Task	7
The Global Shoreline Vector and the New Global Islands Datalayer	
Summary of Method and Results	
Naming the Islands	
Island Maps	9
The Global Island Explorer	22
Accessing the Data	22
Conclusion	23
Acknowledgments	23
References Cited	23

Table 1. Size classes of global islands.

List of Figures

Figure 1. Global islands.

Figure 2. The islands of the Americas.

Figure 3. The islands of Africa and Europe.

Figure 4. The islands of Australia and the southwest Pacific Ocean.

Figure 5. The islands of the Caribbean Sea.

Figure 6. The islands of the Aegean Sea.

Figure 7. The islands of Patagonia

Contents

List of Tables

Figure 8. The islands of New Caledonia.

Figure 9. The islands of Belize.

Figure 10. The islands of the Florida Keys, USA, and northwestern Cuba.

Figure 11. The islands of the Outer Hebrides Islands, Scotland.

Figure 12. The welcome page of the Global Island Explorer, a freely accessible online tool that allows guery and visualization of approximately 341,000 islands.

A New Map of Global Islands

By Roger Sayre, Jack Dangermond, Dawn Wright, Sean Breyer, Kevin Butler, Keith Van Graafeiland, Charles Frye, D. Karagulle, Steve Kopp, Suzanne Noble, Jill Cress, Devon Burton, Madeline Martin, and Jac Steiner.

Abstract

As part of an intergovernmental commission for a high spatial resolution and data-derived global coastal ecosystems map, a new datalayer and map of global islands was produced. The new data represent the most standardized, current, and detailed global islands data yet available. The 340,691 mapped islands were classed by size into 5 continental mainlands, 21,818 big islands (>1km²), and 318,868 small islands (<1km²). The GIS data are available in the public domain as high spatial resolution (30 m) polygons. Herein we 1) summarize the objectives and methods for producing the new islands data, 2) describe the ongoing effort to name the bigger islands (>1km²), 3) present a series of graphics for a visual depiction of the islands at global, regional, national and local scales, 4) present the Global Islands Explorer as an online visualization and query tool developed as a simple window into the data for use by a variety of audiences, and 5) describe the availability of the data.

Introduction

Geographic Data on Global Islands

The surface of the earth can be viewed as primarily an ocean matrix with emergent land areas of varying sizes. Regardless of the size of these land areas, which range from very large landmasses like the Eurasian mainland to tiny coastal rock outcrops, every land area on Earth is ultimately surrounded by ocean waters, and is therefore an island. Islands are increasingly easy to detect in satellite imagery, and even very small ones can be universally explored in broadly accessible technologies like Google Earth®. There are also several list-based compendia of information on islands such as the UNEP Islands Directory (Dahl, 1991) and Wikipedia's Lists of Islands (https:// en.wikipedia.org/wiki/Category:Lists of islands), which are organized by several subcategories including name, size, continent, nation and ocean. Detailed, global characterizations of specific types of islands such as barrier islands (Stutz and Pilkey, 2000) and atoll islands (Goldberg, 2016) are available. However, current, accurate, high resolution digital datasets containing polygonal geospatial representations of all global islands are not generally available in the public domain. We present herein a new, globally comprehensive, high spatial resolution (30 m), GIS dataset of the islands of the earth.

Islands and Biodiversity

Except for the most remote and climatically inhospita-

ble areas on the planet, islands are generally inhabited by a variety of plants and animals, often including humans, if they are big enough to support these organisms. High levels of endemism in flora and fauna are often characteristic of islands due to their geographic isolation (Whittaker and Fernández-Palacios, 2007). This isolation and remoteness represents a barrier to the geographic dispersal of propagules, and over time favors evolutionary adaptations and speciation. Darwin formulated much of his thinking on evolution from his observations of differences in species' traits recorded in his travels to the Galapagos Islands. While species richness on islands is generally regarded as lower than counterpart mainland areas, "endemism richness," an index of species richness scaled by endemism, is relatively high (Kier et al., 2009).

High levels of both endemism and endemism richness on islands suggests that they should be considered as important candidate geographies for biodiversity conservation efforts. This is particularly true given that extinctions and other losses of biodiversity on islands are disproportionally rapid (Spatz et al., 2017). Island species are often vulnerable to human-caused disturbances such as the introduction of invasive species, hunting, and the agriculture-related destruction of relatively small and fragile habitats which contain small populations of rare species (Szabo et al., 2012). A focus of conservation attention on global islands is clearly warranted because 1) their biodiversity importance is high (Kier et al., 2009), and 2) that biodiversity is highly threatened (Spatz et al., 2017). Survey (USGS) is the designated federal agency implementing the work.

Conserving global island biodiversity necessarily re-In response to that commission from GEO, the USGS quires a knowledge of where islands are, what ecosystems and Esri established a public/private partnership and are found on those islands, what condition those ecosysmapped terrestrial ecosystem distributions using a structems are in, and what rare and endangered species are ture-based mapping approach where the ecosystems were found in those ecosystems. To advance understanding of delineated from a vertical integration of the climate regime, the island distributions of globally threatened vertebrates, landforms, substrate, and land cover (Savre et al., 2014). the Threatened Island Biodiversity Database (TIBD, http:// That effort produced a set of global ecological land units, or tib.islandconservation.org/) was developed (Spatz et al., ELUs, as physically distinct areas and their associated veg-2017). The TIBD is a globally comprehensive database etation. Subsequent to that terrestrial ecosystem mapping on 1,189 highly threatened vertebrate species breeding on effort, and again responding to the GEO intergovernmental 1,288 islands. In addition to threatened vertebrate species charge to produce standardized, robust, and practical global and invasive species information, the TIBD provides atecosystem maps for marine environments, the USGS/Esri tribute information on islands such as coordinate location, team then developed a similar method for stratifying the name, and size, and locates islands as points on a world global ocean into physically and chemically distinct volumetric regions. These oceanic, three dimensional, pelagic map. regions are called ecological marine units, or EMUs (Savre A geospatial datalayer which allowed for the visualization, query, and spatially explicit analysis of species et al., 2017a; Sayre et al., 2017b).

and ecosystems' distributions on all islands would con-While the EMUs characterize the distinct chemical and siderably advance understanding of global island biodiphysical environments in the oceanic water column, they versity. Such a layer would permit geospatial analysis of do not adequately identify coastal ecosystems, which are species presence/absence data, population locations, range smaller, often linear, sometimes densely populated ecosizes, and areas of threat, and would support conservation systems at the interface of the land and the sea. The team priority setting within and across islands. As mentioned therefore decided to produce a set of global ecological above, such a geospatial global islands database was not coastal units (ECUs), and developed a methodology for heretofore available, and was produced as part of an inthe approach (Sayre et al., 2018). The work to produce tergovernmental commission. the ECUs is underway.

The GEO (Group on Earth Observation) Global Ecosystem Mapping Task

The Group on Earth Observations (GEO) is a con-One of the prerequisites for the ECU mapping was the development of a new, high spatial resolution, image-desortium of over 100 nations which seeks to leverage the rived global shoreline vector (GSV), which was produced use of Earth observations to help solve some of society's greatest challenges (GEO, 2005). To that purpose, GEO using composite 2014 Landsat satellite imagery at a 30 m has developed an intergovernmental protocol and asresolution (Sayre et al., 2018). The GSV was then used sociated workplan which includes an initiative (GI-14 as the source geospatial linework for the development GEO ECO, http://www.earthobservations.org/activity. of the new global islands datalayer (GID). The GID was php?id=116) on global ecosystems. The initiative fortherefore produced "indirectly," i.e. it was not a primary mally commissions the development of a standardized, objective of the team to produce a new global islands dataset. As an intermediate product associated with the robust, and practical map of global ecosystems for terrestrial, freshwater, and marine environments (Sayre mapping of global coastal ecosystems, the GID resource has considerable additional value beyond characterizing et al., 2007). The United States is the member nation of GEO leading this activity, and the U.S. Geological ecosystems.

The Global Shoreline Vector and the New Global Islands Datalayer

Summary of Method and Results

Detailed methods and results for the production of the GSV and the GID are presented in Savre et al. (2018). In summary, a shoreline vector was extracted from 30 m spatial resolution Landsat imagery from 2014 by visual feature interpretation. An analyst manually "ran" the entire global coastline and interpreted two feature classes, land and ocean, from all of the Landsat images of the planet that contained a segment of shoreline. Antarctica was not included in the analysis due to difficulty in separating ice features from land features. The interpretation was implemented in the cloud (Google Earth Engine®) to minimize the local storage and manipulation of hundreds of Landsat images. Training points collected by the analyst were used to classify the imagery with a semi-automated, statistically rigorous classification algorithm.

A minimum mapping unit of four contiguous 30 m landsat pixels (3600m², or .0036km²) was used, such that any island smaller than this delineation threshold size is not included in the GID datalayer. The classified coastline was subsequently brought down from the cloud, converted from raster to vector format, and cleaned. All segments were joined, and topology was applied to the data to convert line segments to polygons.

Following conversion of coastline segments to island polygons, 340,691 polygons were produced. The five largest polygons were the mainlands of the five continents (North America, South America, Africa, Australia, and Eurasia), with an algorithmic separation of the Americas at the Panama Canal, and a separation of Africa from Eurasia at the Suez Canal. These five polygons were assigned a size class of "continental mainlands." The remaining set of islands greater than 1km² were then attributed as "big islands," and all remaining polygons (<1km²) were attributed as "small islands." Table 1, reproduced from Sayre et al. (2018), shows the number of islands in each of the three size classes, as well as their summed areas and coastline lengths. The 1 km size threshold for separating small and big islands has no ecological basis, and was arbitrarily chosen for ease of understanding and use.

Table 1. The number, size, and coastline length for islands in three size classes. Reproduced from Savre et al. (2018).

Landmass type	Number of polygons	Area (km ²)	Length of coastline (km)
Continental mainlands	5	125,129,046	813,467
Islands > 1 km ²	21,818	9,938,964	1,304,762
Islands $\leq 1 \text{ km}^2$	318,868	20,589	321,774

Naming the Islands

Having derived the 340,691 islands from primary interpretation of satellite imagery, no attribute information on the islands other than the feature counts and geometric information in Table 1 was initially available. The first attribution step was to manually name the largest islands using the UNEP Islands Directory (Dahl, 1991). The 1,000 largest islands on the UNEP list were located in the GID, and their names were manually added to the feature attribute table. Three sources of information were consulted when searching for names for islands: Esri's

WorldImage Basemap®, Open Street Map®, and Google Maps[®]. This manual naming process was considered accurate, but was quite time-intensive. When naming the largest 1000 islands, an additional ~2000 islands were opportunistically named (i.e. when zoomed in to name one of the 1000 largest islands it was often expedient to name several islands in the immediate vicinity at that zoom level). At the end of this first phase of manual naming, 3,156 big islands (>1km²) were named, and 686 small islands (<1km²) were named.

To facilitate and hasten the naming, an automated approach was then adopted. The GID was intersected with the GeoNames server. GeoNames is a geographical names sateague," not "Assateague Island." This convention was database freely available in the public domain containing sometimes difficult to enforce with islands with non-Enover eleven million point-based placenames for a variety glish names, and exceptions to the convention were made. of features, including islands (https://www.geonames. For example, there is an island in the Dominican Republic org/). The GeoNames island placename (any placename called Isla Saona. An English equivalent would be Saona with an ISL feature class) that was closest to the centroid Island, which would then be attributed simply as Saona. of an island polygon was added as the name attribute to However, the island is universally referred to as Isla Saona, the GID. The placename was only used if it was located by English speakers and non-English speakers alike. The inside the island polygon. Any island that had previously name was therefore attributed as Isla Saona. Whenever been manually named retained that name. The reconcilpossible, however, the name attribute was limited only to iation of the GeoNames database with the GID resulted the proper noun. in the addition of 5,340 automated big island names, and Only the 26 letters of the English alphabet (ISO basic 12,253 automated small island names. The total number Latin alphabet) were used. For simplification, diacritical of named islands at the conclusion of the automated phase marks (accents, cedillas, tildes, umlauts, etc.) were elimwas 12,939 (0.04 %) small islands and 8,496 (38.9 %) big inated from placenames, such that all names have been islands. The automated naming approach was successful reduced to the simplest possible English language formuin retrieving and assigning thousands of names, but more lation. The attribute "UNNAMED" was given to islands than 60 percent of the big islands and 99 percent of the when either a name was not found, or a name was available small islands still lacked names. but not with English language characters. In the latter case, it is understood that the attribute UNNAMED more At this point in the process, a decision was made to accurately means the name was not available in English.

discontinue the naming effort for the small islands, due to the enormity of the effort that would be required to name Many of the islands identified by the classification althe approximately 305,000 remaining small islands, many gorithm were fluvial islands or islands near strongly disof which were likely small outcrops that lacked a name. sected coastlines. In these deltaic systems, where islands Work is now underway to complete the naming process are dynamic and island shapes are often changing, the for the remaining approximately 13,000 big islands with a islands often lack names in the source data. In these cases, islands were left unnamed, or were in some cases named return to the labor intensive manual naming method, again using Esri's WorldImage Basemap®, Open Street Maps®, according to the river in which they were located. For and Google Maps[®] as the source datalayers for the names. example, several unnamed islands in the mouth of the A number of conventions were used in the naming Orinoco River were given an attribute name of "Orinoco process. Only the proper noun, in English, was included in Delta." Other islands, for example in coastal or mangrove habitats, that lacked a name in the source data were simply the attribute field. For example, the polygon corresponding to Assateague Island was given a name attribute of "Asattributed as UNNAMED, as described above.

A sequence of graphics is presented depicting islands exploration of the global islands data, and demonstrate both what is discernible and the level of linework detail at global (Figure 1), continental (Figure 2, the Americas; Figure 3, Africa and Europe; Figure 4, Australia and the at resolutions ranging from very coarse to very fine. At southwest Pacific Ocean), regional (Figure 5, the Caribcoarse resolutions and small page formats, small islands bean Sea; Figure 6, the Aegean Sea; Figure 7, Patagonia) are generally not visible. They appear with increasing national (Figure 8, New Caledonia; Figure 9, Belize), and detail as the resolution is increased. These maps are insubnational (Figure 10, Florida Keys; Figure 11, Outer tended to provide visual impressions of the numbers and variety of shapes and sizes of Earth's islands. Hebrides) scales. These maps are presented as a visual

Island Maps



Figure 1. Global islands. All islands except continental mainlands are depicted in a geen color. Small islands, not visible at this coarse resolution and page format size, have been rendered visible through visual exaggeration using increased point sizes for feature symbols.

Figure 3. The islands of Africa and Europe.





Figure 4. The islands of Australia and the southwest Pacific Ocean.





Figure 5. The islands of the Caribbean Sea. Large islands (>1km²) are depicted in green; small islands (<1km²) are depicted in gold.









Figure 8. The islands of New Caledonia. Large islands (>1km²) are depicted in green; small islands (<1km²) are depicted in gold.

Figure 9. The islands of Belize. Large islands (>1km²) are depicted in green; small islands (<1km²) are depicted in gold.





A Special Publication of the American Association of Geographers

Figure 10. The islands of the Florida Keys, USA, and northwestern Cuba. Large islands (>1km²) are depicted in green; small islands (<1km²) are depicted in gold.

· · · · 3

Figure 11. The islands of the Outer Hebrides Islands, Scotland. Large islands (>1km²) are depicted in green; small islands (<1km²) are depicted in gold.



The Global Island Explorer

The global islands data can easily be visualized, queried and analyzed using GIS technologies, but to make the data easily accessible to the broadest possible audience we also developed a Global Islands Explorer tool (GIE: https:// rmgsc.cr.usgs.gov/gie/). The GIE is a web-based island data exploration application that only requires internet access for use. It is an open access resource which will be maintained, and GIE functionality will eventually be expanded

beyond simple exploration to include limited analysis (e.g. feature counts, buffering, etc.) Currently, pan, zoom and query functions are included, as is a text-based exploration tool allowing users to explore any geography of interest by typing in the name of an island or region. The island data are served as raster image services, but are available for download in their original vector polygon format. The welcome page for the GIE is presented in Figure 12.

Accessing the Data

The global islands data and the GIE tool are open data resources available without a login requirement from the Esri Living Atlas (http://livingatlas.arcgis.com) and elsewhere in the ArcGIS Online content. The islands data are therefore easily integrated with hundreds of other datasets representing the most current, detailed, authoritative, and curated GIS-ready global islands data available. This integration is easily accomplished without the need for

downloading, preparing, and reconciling disparate datasets by the user. The islands data are also available in the public domain at the U.S. Geological Survey's Global Ecosystems webpage and resources (https://rmgsc.cr.usgs.gov/ outgoing/ecosystems/Global/). While the data are available in ArcGIS formats, they are also accessible as polygon features in an Open Geospatial Consortium (OGC) Geo Package, or similar, format.

Figure 12. The welcome page of the Global Island Explorer tool.



Global Islands Data

Reference:

Sayre, R., S. Noble, S. Hamann, R. Smith, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, C. Frye, D. Karagulle, D. Hopkins, D. Stephens, K. Kelly, Z, basher, D. Burton, J. Cress, K. Atkins, D. van Sistine, B. Friesen, B. Allee, T. Allee, P. Aniello, I Asaad, M. Costello, K. Goodin, P. Harris, M. Kavanaugh, H. Lillis, E. Manca, F. Muller-Karger, B. Nyberg, R. Parsons, J. Saarinen, J. Steiner, and A. Reed. 2018. A new 30 meter resolution global shoreline vector and associated global islands database for the development of standardized global ecological coastal units. Journal of Operational Oceanography – A Special Blue Planet Edition. DOI:10.1080/1755876X.2018.1529714. https://doi.org/10.1080/1755876X.2018.1529714

Contacts:

toger Sayre, U.S. Geological Survey

GIE Developer and Webmaster Cress, U.S. Geological S

rhvacy Policy | Legal | Accessibility | Site Map | Contact USG

In a first-of-its kind effort, spatial data on the location and shapes of the islands of the Earth were extracted from satellite imagery in a standardized and replicable proces A new global islands datalayer was produced which more current, comprehensive, and accurate than existin global island data products. The new islands data are avail able as polygons for use in GIS technologies, and for othe non-specialist audiences the data are easily accessed an

Acknowledgments

We acknowledge and thank the American Association of Geographers, and Doug Richardson in particular, f support in producing and promoting this and simil work at several AAG Annual Meetings over the pa ten years. We thank Becky Pendergast for extraordina design and layout support in all of the booklets in the

- Dahl, A.1991. Island directory. UNEP Regional Seas Directories an Bibliographies Number 35. Nairobi, UN Environment progra 573 pages.
- GEO (Group on Earth Observations). 2005. Global Earth Observ tion Systems (GEOSS) 10 Year Implementation Pl Reference Document. ESA Publication GEO 1000R/ESA SP 128 Noordwijk, The Netherlands, European Space Agency. 210 pag
- Goldberg, W. 2016. The Atolls of the World Revisiting the Origin Checklist. Atoll Research Bulletin Number 610. Washington, De Smithsonian Instituition Scholarly Press. 47 pages. DOI: 10.547 si.0077-5630.610
- Kier, G., H. Kreft, T. Lee, W. jetz, P. Ibisch, C. Nowicki, J. Mutke, a W. Barthlott. 2009. A global assessment of endemism and spec richness across island and mainland regions. Proceedings of t National Academy of Sciences 106(23): 9322-9327. www.pna org cgi doi 10.1073 pnas.0810306106
- Savre, R., D. Muchoney, and A. Yanosky. 2007. Mapping glob ecosystems - The GEOSS approach. Pages 231-232 in The F Picture, Geneva, Switzerland: GEO, 278 pages.
- Sayre, R., J. Dangermond, C. Frye, R. Vaughan, P. Aniello, S. Brey D. Cribbs, D. Hopkins, R. Nauman, W. Derrenbacher, D. Wrigh C. Brown, C. Convis, J. Smith, L. Benson, D. Paco VanSistine, Warner, J. Cress, J. Danielson, S. Hamann, T. Cecere, A. Reddy, Burton, A. Grosse, D. True, M. Metzger, J. Hartmann, N. Moosdo H. Dürr, M. Paganini, P. DeFourny, O. Arino, S. Maynard, Anderson, and P. Comer. 2014. A New Map of Global Ecolog cal Land Units - An Ecophysiographic Stratification Approa Washington, DC: Association of American Geographers. 46 pag
- Sayre, R, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, Costello, P. Harris, K. Goodin, J. Guinotte, Z. Basher, M. Kavar ugh, P. Halpin, M. Monaco, N. Cressie, P. Aniello, C. Frye, and Stephens. 2017a. A three-dimensional mapping of the ocean bas on environmental data. Oceanography. 30(1):90-103. doi:10.567 oceanog.2017.116.

Conclusion

ns	explored using a new web-based Global Island Explorer
m	tool. The data are intended to be useful in assessments of
SS.	island biodiversity, ecological processes, and ecosystem
is	accounting, and could also be useful for resource manage-
ng	ment and planning at multiple scales. Maps describing the
il-	island distributions at global, continental, regonal, national
er	and subnational scales are presented herein. Access to the
nd	data is open.

on	series, of which this work is the fourth. We thank Jesse
or	Wong and Yufen Huang, affiliated with the U.S. Geolog-
ar	ical Survey, for helpful reviews of the manuscript. Any
ıst	use of trade, product, or firm names is for descriptive
ry	purposes only, and does not imply endorsement by the
nis	U.S. Government.

References Cited

nd m. va-	Sayre, R., J. Dangermond, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, M. Costello, P. Harris, K. Goodin, M. Kavanaugh, N. Cressie, J. Guinotte, Z. Basher, P. Halpin, M. Monaco, P. Aniello, C. Frye, D. Stephens, P. Valentine, J. Smith, R. Smith, D. VanSistine, L. Course, M. Wasser, G. Davier, J. Stafford, D. C. Karlow, D. Vanser, C. Davier, J. Stafford, D. C. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Smith, S. Sayre, D. Stephens, P. Valentine, J. Smith, R. Smith, D. Vanser, J. Smith, S. Sayre, D. Stephens, P. Valentine, J. Smith, S. Sayre, D. Stephens, P. St
lan 84. es. nal	J. Cress, H. Warner, C. Brown, J. Sterrenson, D. Cribos, B. Van Esch, D. Hopkins, G. Noll, S. Kopp, and C. Convis. 2017b. A New Map of Global Ecological Marine Units – An Envrionmental Stratification Approach. Washington, DC: Association of American Geographers. 35 pages.
nd ies the as. pal ull	 Sayre, R., S. Noble, S. Hamann, R. Smith, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, C. Frye, D. Karagulle, D. Hopkins, D. Stephens, K. Kelly, Z. Basher, D. Burton, J. Cress, K. Atkins, D. Van Sistine, B. Friesen, R. Allee, T. Allen, P. Aniello, I. Asaad, M. Costello, K. Goodin, P.Harris, M. Kavanaugh, H. Lillis, E. Manca, F. Muller-Karger, B. Nyberg, R. Parsons, J. Saarinen, J. Steiner, and A. Reed. 2018. A new 30 meter resolution global shoreline vector and associated global islands database for the development of standardized ecological coastal units, Journal of Operational Oceanography, DOI:10.1080/1755876X.2018.1529714
er, ht, H.	Spatz, D. K. Zilliacus, N. Holmes, S. Butchart, P. Genovesi, G. Ceballos, B. Tershy, and D. Croll. 2017. Globally threatedned vertebrates on islands with invasive species. Science Advances, 3(10):e1603080. DOI: 10.1126/sciadv.1603080
D. orf, M.	Stutz, M., and O. Pilkey. 2000. A review of global barrier island distri- bution. Journal of Coastal Research Special Issue 34. Proceedings ICS 2000, New Zealand. ISSN 0749-0208, Pages 15-22.
gi- ch. es. M.	Szabo, J., S. Butchart, H. Possingham, and S. Garnet. 2012. Adapting global biodiversity indicators to the national scale: A Red List Index for Australian birds. Biological Conservation, 148: 61-68, DOI: 10.1016/j.biocon.2012.01.062
D. sed 70/	Whittaker R., and J. Fernandez-Palacios. 2007. Island Biogeography: Ecology, Evolution, and Conservation. 2nd edn. Oxford University Press, Oxford. 416 pages.